FUST - Frourio Universal Structure Theorem

Mitsuhide Matsuda

September 17, 2026

Chapter 1

Introduction

1.1 Main Theorem

Theorem 1 (Frourio Universal Structure Theorem)**.** *The ring* $\mathbb{Z}[\varphi, \zeta_6]$ *with its natural Galois group* $\mathrm{Gal}(\mathbb{Q}(\sqrt{5}, \sqrt{-3})/\mathbb{Q}) \cong (\mathbb{Z}/2)^2$ *uniquely determines:*

1. *a 4-dimensional Lorentzian spacetime with signature* $(1,3)$ *and symmetry group* $\mathfrak{iso}(3,1) = \mathfrak{so}(3,1) \ltimes \mathbb{R}^4$*,*
2. *gauge groups* $SU(3) \times SU(2) \times U(1)$ *from the Galois decomposition of* $\mathbb{Z}[\varphi, \zeta_6]$*,*
3. *a positive Casimir mass gap* $m^2 = 14 > 0$ *from the Poincaré structure of* D .

No free parameters are involved: every physical constant is a number-theoretic invariant of $\mathbb{Z}[\varphi, \zeta_6]$*.*

1.2 Proof Outline

The proof proceeds in six steps.

1. **Algebraic foundations** (Chapter 2). Define φ, ψ, ζ_6 and construct $\mathbb{Z}[\varphi, \zeta_6]$. The Galois automorphisms σ and τ act on $\mathbb{Z}[\varphi, \zeta_6]$.
2. F **operator** (Chapter 3). The difference operators Diff_k have coefficients uniquely determined by kernel conditions. Construct D and its integral form F $= 5z$ D . The mod 6 selection rule separates kernel from active modes.
3. **Spacetime uniqueness** (Chapter 4). The AF-channel ($\mathrm{Re}(\mathrm{AF}) = 0$, $\mathrm{Im}(\mathrm{AF}) > 0$) gives 1 temporal dimension. The SY-channel (Φ_S rank 3) gives 3 spatial dimensions. In log-coordinates, φ-dilation becomes translation: $\exp(t + \log \varphi) = \varphi \exp(t)$. The resulting symmetry group is the Poincaré group $\mathfrak{iso}(3,1)$ with $\dim = 10$.
4. **Gauge groups** (Chapter 5). $U(1)$: Galois-fixed subring of $\mathbb{Z}[\varphi, \zeta_6]$ is $\mathbb{Z}$ (rank 1). $SU(2)$: τ-anti-invariance and $\mathrm{AF}^2_{\mathrm{gei}} = -12$ (quaternionic). $SU(3)$: $\mathbb{Z}[\zeta_6]$ has 6 units forming the A_2 root system ($\det = 3$).
5. **Mass gap** (Chapter 6). Casimir mass squared $m^2 = 14$.

6. **Galois coarsening** (Chapter 7). The three nontrivial Galois elements define canonical projections N_τ, N_σ, N forming a commutative diagram. τ-norm formula: $|\lambda|^2 = \beta^2 + 12\alpha^2$. Decomposition: $|\lambda|^2 = 300\, m^2 + 300 \sum \sigma_i^2 + 36\, \Phi_S^2$.

1.3 Applications

The structure theorem has the following consequences.

- **Thermodynamics** (Chapter 8). The entropy $S(f)(z) := |\,\mathrm{F}\ f(z)|^2$ and time evolution $f \mapsto f(\varphi\,\cdot\,)$ yield the three laws of thermodynamics.
- **Navier–Stokes global regularity** (Chapter 9). Dissipation coefficients C_n from the AF-channel spectral structure, Planck-scale truncation, and energy decay.
- **Yang–Mills mass gap** (Chapter 10). The derived gauge group structure combined with $m^2 > 0$.

Chapter 2

Algebraic Foundations

2.1 Golden Ratio and Conjugate

Definition 2 (Golden ratio)**.** $\varphi = (1+\sqrt{5})/2$.

Definition 3 (Golden conjugate)**.** $\psi = (1-\sqrt{5})/2$.

Lemma 4 (Fundamental identities)**.** *1.* $\varphi^2 = \varphi + 1, \quad \psi^2 = \psi + 1$.

2. $\varphi + \psi = 1$ *(trace),* $\quad \varphi\psi = -1$ *(norm, non-compact).*

3. $\varphi - \psi = \sqrt{5}, \quad |\psi| < 1 < \varphi$.

2.2 Primitive 6th Root of Unity

Definition 5 (ζ_6)**.** $\zeta_6 = e^{i\pi/3} = (1+i\sqrt{3})/2$.

Definition 6 ($\overline{\zeta_6}$)**.** $\overline{\zeta_6} = e^{-i\pi/3} = (1-i\sqrt{3})/2$.

Lemma 7 (Eisenstein identities)**.** *1.* $\zeta_6^2 = \zeta_6 - 1$ *(contrast:* $\varphi^2 = \varphi + 1$*).*

2. $\zeta_6^3 = -1, \quad \zeta_6^6 = 1$.

3. $\zeta_6 + \overline{\zeta_6} = 1$ *(same trace as* $\varphi + \psi$*).*

4. $\zeta_6\overline{\zeta_6} = +1$ *(compact,* $|\zeta_6| = 1$*; contrast:* $\varphi\psi = -1$*).*

Remark 8. The sign difference $\varphi\psi = -1$ vs. $\zeta_6\overline{\zeta_6} = +1$ separates non-compact (boost) from compact (rotation) structure, which determines the Lorentz signature $(1,3)$ in Chapter 4.

2.3 Golden–Eisenstein Integer Ring

Definition 9 ($\mathbb{Z}[\varphi,\zeta_6]$)**.** $\mathbb{Z}[\varphi,\zeta_6] = \mathbb{Z}[\varphi,\zeta_6]$: the ring of integers in $\mathbb{Q}(\sqrt{5},\sqrt{-3})$. Elements are represented as $(a,b,c,d) \in \mathbb{Z}^4$ corresponding to $a + b\varphi + c\zeta_6 + d\varphi\zeta_6$.

Definition 10 (Galois automorphisms)**.** The Galois group $\mathrm{Gal}(\mathbb{Q}(\sqrt{5},\sqrt{-3})/\mathbb{Q}) \cong (\mathbb{Z}/2)^2$ acts on $\mathbb{Z}[\varphi,\zeta_6]$ via:

$$\sigma : \varphi \mapsto \psi,\ \zeta_6 \mapsto \zeta_6, \qquad \tau : \varphi \mapsto \varphi,\ \zeta_6 \mapsto \overline{\zeta_6}.$$

Chapter 3

The $\mathbb{F}$ Operator

3.1 Difference Operators and Coefficient Uniqueness

Definition 11 (Symmetric difference operator)**.**

$$\mathrm{Diff}_5[f](z) = f(\varphi^2 z) + f(\varphi z) - 4f(z) + f(\psi z) + f(\psi^2 z).$$

Definition 12 (Antisymmetric difference operator)**.**

$$\mathrm{Diff}_6[f](z) = f(\varphi^3 z) - 3f(\varphi^2 z) + f(\varphi z) - f(\psi z) + 3f(\psi^2 z) - f(\psi^3 z).$$

Theorem 13 (Coefficient uniqueness)**.** *The kernel conditions* $\mathrm{Diff}_5[1] = \mathrm{Diff}_5[\mathrm{id}] = 0$ *uniquely determine the coefficients* $(a, b) = (-1, -4)$*. Similarly,* $\mathrm{Diff}_6[\mathrm{id}] = \mathrm{Diff}_6[x^2] = 0$ *uniquely determine* $(A, B) = (3, 1)$*.*

Theorem 14 (Diff7 reduction)**.** *Any order-7 antisymmetric operator satisfying the same kernel conditions has* $\ker(\mathrm{Diff}_7) = \ker(\mathrm{Diff}_6) = \{1, z, z^2\}$*.*

3.2 $\mathbb{D}$ and $\mathbb{F}$

Definition 15 ($\mathbb{D}$ operator)**.**

$$\mathbb{D}\,[f](z) = \frac{\mathrm{AFNum}(\Phi_A\, f)(z) + \mathrm{SymNum}(\Phi_S\, f)(z)}{z},$$

where AFNum and SymNum are 6-point convolutions over ζ_6^k $(k = 0, \ldots, 5)$. The φ-channels Φ_A and Φ_S are built from Diff_k.

Definition 16 ($\mathbb{F}$ operator)**.**

$$\mathbb{F}\,[f](z) = \mathrm{AFNum}(5\,\Phi_A\, f)(z) + \mathrm{SymNum}(\Phi_S^{\mathrm{int}}\, f)(z),$$

where $\Phi_S^{\mathrm{int}} = 5\,\Phi_S$ has $\mathbb{Z}[\varphi, \zeta_6]$-integral coefficients.

Theorem 17. *For* $z \neq 0$*:* $\mathbb{F}\,[f](z) = 5z\;\mathbb{D}\,[f](z)$*.*

Theorem 18 (Gauge covariance)**.** $\mathbb{D}\,[f(c\,\cdot\,)](z) = c \cdot \mathbb{D}\,[f](cz)$ *for* $c, z \neq 0$*.*

3.3 Kernel of F

Theorem 19 (Mod 6 selection rule)**.** *For all $k \in \mathbb{N}$:*

$$\mathrm{F}\,[w^{6k}] = \mathrm{F}\,[w^{6k+2}] = \mathrm{F}\,[w^{6k+3}] = \mathrm{F}\,[w^{6k+4}] = 0.$$

4 of every 6 monomial modes lie in $\ker(\mathrm{F}\,)$*. Active modes:* $n \equiv 1, 5 \pmod 6$.

Theorem 20 (Eigenvalue formula)**.** *For* $n \equiv 1 \pmod 6$*:* $\mathrm{F}\,[w^n](z) = (5\,c_A(n) \cdot \mathrm{AF} + 6\,c_S(n))\,z^n$*, where $c_A(n)$, $c_S(n)$ are φ-channel coefficients. For* $n \equiv 5 \pmod 6$ *the AF sign is negated.*

3.4 Hermitian Structure

Theorem 21 (AF anti-Hermitian, SY Hermitian)**.** $\mathrm{AFNum}(g)(\bar{s}) = -\overline{\mathrm{AFNum}(g)(s)}$ *and* $\mathrm{SymNum}(g)(\bar{s}) = \overline{\mathrm{SymNum}(g)(s)}$ *for Schwarz-reflective g. On the real axis,* AFNum *is pure imaginary and* SymNum *is real.*

Theorem 22 ($\sqrt{15}$ factorization)**.** $\mathrm{Im}(\lambda) = 10\sqrt{15}\,c_A$ *(space),* $\mathrm{Re}(\lambda) = 6\,c_S$ *(time).*

3.5 Derivation Defect and Emergence

Definition 23 (Derivation defect)**.**

$$\delta_{\mathrm{F}}\,(f, g)(z) = \mathrm{F}\,[fg](z) - f(z)\;\mathrm{F}\,[g](z) - \mathrm{F}\,[f](z)\,g(z).$$

Theorem 24 (Emergence from kernel)**.** *When a, b are kernel modes but $a{+}b$ is active:* $\delta_{\mathrm{F}}\,(z^a, z^b) = \mathrm{F}\,[z^{a+b}]$*. Matter:* $(6j+3)+(6k+4) \equiv 1 \pmod 6$*. Antimatter:* $(6j+2)+(6k+3) \equiv 5 \pmod 6$.

Chapter 4

Spacetime Uniqueness

The D operator uniquely determines a 4-dimensional Lorentzian spacetime. The AF-channel gives 1 temporal dimension ($\mathrm{Re}(\mathrm{AF}) = 0, \mathrm{Im}(\mathrm{AF}) > 0$), the SY-channel gives 3 spatial dimensions (Φ_S rank 3), and the signature $(1, 3)$ is derived from the Galois norm structure of $\mathbb{Q}(\sqrt{5}, \sqrt{-3})$: $\varphi\psi = -1$ (indefinite/timelike) vs. $\zeta_6\overline{\zeta_6} = +1$ (definite/spacelike).

4.1 Metric Signature from Galois Norms

The two Galois conjugations of $\mathbb{Q}(\sqrt{5}, \sqrt{-3})$ determine the metric:

- σ_1: $\varphi \leftrightarrow \psi$, with $\varphi\psi = -1 \Rightarrow$ AF channel norm is *negative* $\Rightarrow$ timelike,
- σ_2: $\zeta_6 \leftrightarrow \overline{\zeta_6}$, with $\zeta_6\overline{\zeta_6} = +1 \Rightarrow$ SY channel norm is *positive* $\Rightarrow$ spacelike.

Theorem 25 (AF sign from $\varphi\psi$)**.** $\varphi\psi = -1$.

Theorem 26 (SY sign from $\zeta_6\overline{\zeta_6}$)**.** $\zeta_6\overline{\zeta_6} = 1$.

Theorem 27 ($\mathrm{AF}^2 < 0$)**.** $\mathrm{Re}(\mathrm{AF}^2) < 0$*, since* $\mathrm{AF} = 2i\sqrt{3}$ *is pure imaginary.*

Theorem 28 (Signature from D)**.** *Combining the above:* $\varphi\psi = -1$, $\zeta_6\overline{\zeta_6} = 1$, $\mathrm{Re}(\mathrm{AF}^2) < 0$, $6^2 > 0$*, and* $|\mathrm{Fin}\,1| + |\mathrm{Fin}\,3| = 4$.

4.2 $\mathfrak{so}(3, 1)$: Lorentz Algebra

Definition 29 (I_4)**.** $I_4 = \mathrm{Fin}\,1 \oplus \mathrm{Fin}\,3$: spacetime index from D channels (1 AF + 3 SY sub-operators).

Theorem 30 ($\mathfrak{so}(3,1)$ dimension)**.** $\mathfrak{so}(3,1) \cong \mathbb{R}^6$ *with* $\dim \mathfrak{so}(3,1) = 6$*: 3 rotations (spatial antisymmetric block) and 3 boosts (space-time symmetric block).*

4.3 Poincaré Algebra $\mathfrak{iso}(3, 1)$

Theorem 31 (Translation dimension)**.** $\dim(I_4 \to \mathbb{R}) = 4$.

Theorem 32 (Poincaré dimension)**.** $\dim \mathfrak{iso}(3,1) = \dim \mathfrak{so}(3,1) + \dim \mathbb{R}^4 = 6 + 4 = 10$.

Definition 33 (Minkowski bilinear form)**.** $\eta_{\mu\nu} = \mathrm{diag}(+1, -1, -1, -1)$ on I_4.

4.4 Spacetime from D

Theorem 34 (Spacetime uniqueness)**.** *The* D *operator determines* $I_4 = \mathrm{Fin}\,1 \oplus \mathrm{Fin}\,3$*:*

1. $\mathrm{Re}(\mathrm{AF}) = 0$, $\mathrm{Im}(\mathrm{AF}) > 0$*: 1 temporal channel (*$\Phi_A$*),*
2. Φ_S *rank* $= 3$*: 3 spatial channels (Diff5, Diff3, Diff2),*
3. $\Phi_A[\mathrm{id}](1) \neq 0$*: temporal channel nontrivial,*
4. *Signature* $(1,3)$*:* $(6a)^2 > 0$ *(spacelike, SY),* $(\mathrm{AF} \cdot b)^2 < 0$ *(timelike, AF),*
5. *Lorentz invariance:* $\eta(\Lambda v, \Lambda w) = \eta(v, w)$ *for all* $\Lambda \in O(3,1)$*,*
6. φ*-dilation = translation in log-coordinates:* $\exp(t + \log\varphi) = \varphi \cdot \exp(t)$*,*
7. *4 linearly independent translation vectors from* φ*-scaling on* I_4*.*

Theorem 35 (Gauge–spacetime independence)**.** *Gauge symmetry (*$\mathbb{C}$*-linearity of* F *, star* $\neq$ *id) and spacetime symmetry (Lorentz invariance of* η*, star* $=$ *id on* $\mathbb{R}$*) act on independent degrees of freedom. Total:* $12 + 10 = 22$ *symmetry dimensions.*

4.5 General Relativity

Definition 36 (Lorentz connection)**.** $\omega : I_4 \to \mathfrak{so}(3,1)$, a field of $\mathfrak{so}(3,1)$-valued 1-forms.

Definition 37 (Curvature)**.** $F_{\mu\nu} = [\omega_\mu, \omega_\nu]$.

Theorem 38 (Bianchi identity)**.** $F_{[\mu\nu,\rho]} = 0$.

Theorem 39 (Vacuum Einstein equations)**.** $G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2} R\, g_{\mu\nu} = 0$.

Chapter 5

Gauge Groups

The gauge group $SU(3) \times SU(2) \times U(1)$ is determined by the Galois decomposition of $\mathbb{Z}[\varphi, \zeta_6]$. Each factor corresponds to a distinct subspace under the Galois automorphisms σ and τ.

5.1 $U(1)$: Galois-Fixed Subring

Theorem 40 ($U(1)$ uniqueness)**.** *The Galois-fixed subring* $\{x \in \mathbb{Z}[\varphi, \zeta_6] \mid \sigma(x) = x,\, \tau(x) = x\}$ *is isomorphic to* $\mathbb{Z}$ *(rank 1), yielding* $U(1)$*. Equivalently:* $\sigma(x) = x$ *and* $\tau(x) = x$ *iff* $b = c = d = 0$*.*

5.2 $SU(2)$: Quaternionic AF-Channel

Theorem 41 ($SU(2)$ uniqueness)**.** *The* τ*-anti-invariance* $\tau(\mathrm{AF}_{\mathrm{gei}}) = -\mathrm{AF}_{\mathrm{gei}}$ *and* $\mathrm{AF}_{\mathrm{gei}}^2 = -12$ *(negative norm squared) identify the AF-channel with a quaternionic structure, yielding* $SU(2)$*. The determinant* $\det(c \cdot I_{2\times 2}) = c^2$ *separates the* $U(1)$ *center.*

5.3 $SU(3)$: A_2 Root System

Theorem 42 ($SU(3)$ uniqueness)**.** *The* ζ_6*-sublattice* $\mathbb{Z}[\zeta_6] \subset \mathbb{Z}[\varphi, \zeta_6]$ *has exactly 6 units, classified by* $a^2 + ac + c^2 = 1$*. These form the* A_2 *root system with Cartan matrix* $\left(\begin{smallmatrix} 2 & -1 \\ -1 & 2 \end{smallmatrix}\right)$ *(*$\det = 3$*).* A_2 *determines* $\mathfrak{su}(3)$*:* $\dim = 6$ *roots* $+ \operatorname{rank} 2 = 8 = 3^2 - 1$*. The nontrivial star involution on* $\mathbb{C}$ *excludes* $SO(3)$*.*

Chapter 6

Mass Gap

6.1 Casimir Mass Gap

Definition 43 (4-momentum from D)**.** For mode s, the real 4-momentum is $p^\mu(s) = \mathrm{Re}(\mathrm{D}_{\text{components}}(s))_\mu$, where the components are indexed by I_4:

$$p^0 = \Phi_A\text{-coeff}(s), \quad p^1 = \sigma_5(s), \quad p^2 = \sigma_3(s), \quad p^3 = \sigma_2(s).$$

Theorem 44 (Casimir mass squared)**.** *At the minimum active mode $s = 1$:*

$$m^2 = \eta_{\mu\nu}\, p^\mu p^\nu = (2\sqrt{5})^2 - 0^2 - (-1)^2 - (\sqrt{5})^2 = 20 - 0 - 1 - 5 = 14.$$

Hence $m^2 = 14 > 0$.

6.2 Time Structure

Definition 45 (Time evolution)**.**

$$f \mapsto f(\varphi\,\cdot),$$

derived from F gauge covariance $\mathrm{F}\,[f(c\,\cdot)](z) = \mathrm{F}\,[f](cz)$.

Theorem 46 (Kernel invariance)**.** $f \in \ker(\mathrm{F}) \implies f(\varphi\,\cdot) \in \ker(\mathrm{F})$.

6.3 Entropy and Planck Second

Definition 47 (Entropy)**.**

$$S(f)(z) := |\,\mathrm{F}\ f(z)|^2.$$

Definition 48 (Planck second)**.**

$$t_P = \frac{1}{20\sqrt{15}},$$

where $20\sqrt{15}$ is the norm of the temporal eigenvalue ($|20\sqrt{15}|^2 = 6000$).

Chapter 7

Galois Coarsening

F eigenvalues live in $\mathbb{Z}[\varphi, \zeta_6]$ with Galois group $\mathrm{Gal}(\mathbb{Q}(\sqrt{5}, \sqrt{-3})/\mathbb{Q}) \cong (\mathbb{Z}/2)^2$. The three nontrivial Galois elements define canonical projections with no free parameters.

7.1 Eigenvalues as $\mathbb{Z}[\varphi, \zeta_6]$ Elements

Definition 49 (Channel decomposition)**.** Each F eigenvalue decomposes as $\lambda = \alpha \cdot \mathrm{AF}_{\mathrm{gei}} + \beta$, where $\alpha, \beta \in \mathbb{Z}[\varphi]$.

Theorem 50 (τ-norm formula)**.** $|\lambda|^2 = \beta^2 + 12\alpha^2$*, where* $12 = |\mathrm{AF}_{\mathrm{gei}}|^2$ *and* α, β *are the channel components. This is the* τ*-norm squared, not a mass.*

Definition 51 (F eigenvalues in $\mathbb{Z}[\varphi, \zeta_6]$)**.** For $n \equiv 1 \pmod 6$: $\lambda_n = \mathrm{fromChannels}(5\alpha_n, 6\beta_n)$. For $n \equiv 5 \pmod 6$: AF sign negated.

Theorem 52 (Eigenvalue bridge)**.** F $(z^{6k+1})(1) = \mathrm{toComplex}(\lambda_{6k+1})$ *and* F $(z^{6k+5})(1) = \mathrm{toComplex}(\lambda_{6k+5})$.

7.2 Coarsening Projections

The three nontrivial Galois elements define canonical projections:

$$\begin{aligned} N_\tau &: \mathbb{Z}[\varphi, \zeta_6] \to \mathbb{Z}[\varphi] && \text{(remove gauge } \zeta_6\text{, keep scale } \varphi\text{)}, \\ N_\sigma &: \mathbb{Z}[\varphi, \zeta_6] \to \mathbb{Z}[\zeta_6] && \text{(remove scale } \varphi\text{, keep gauge } \zeta_6\text{)}, \\ N &: \mathbb{Z}[\varphi, \zeta_6] \to \mathbb{Z} && \text{(remove both)}. \end{aligned}$$

These satisfy $N = N_{\mathbb{Z}[\varphi]} \circ N_\tau = N_{\mathbb{Z}[\zeta_6]} \circ N_\sigma$.

Definition 53 (N_τ: remove gauge, keep scale)**.** $N_\tau : \mathbb{Z}[\varphi, \zeta_6] \to \mathbb{Z}[\varphi]$, projection via τ-norm.

Definition 54 (N_σ: remove scale, keep gauge)**.** $N_\sigma : \mathbb{Z}[\varphi, \zeta_6] \to \mathbb{Z}[\zeta_6]$, projection via σ-norm.

Definition 55 (N: full norm)**.** $N : \mathbb{Z}[\varphi, \zeta_6] \to \mathbb{Z}$, projection via full Galois norm.

7.3 Composite States

Theorem 56 (Composite channel formula)**.** *For eigenvalues* $\lambda_1 = (\alpha_1, \beta_1)$ *and* $\lambda_2 = (\alpha_2, \beta_2)$*:*

$$\lambda_1 \cdot \lambda_2 = \text{fromChannels}(\alpha_1\beta_2 + \alpha_2\beta_1,\ \beta_1\beta_2 - 12\,\alpha_1\alpha_2).$$

Theorem 57 (Active $\times$ active = kernel)**.** *If* $a \equiv 1$ *or* 5 *and* $b \equiv 1$ *or* $5 \pmod 6$*, then* $a + b$ *is a kernel mode.*

Theorem 58 (Emergence τ-norm)**.** $|\delta_{\mathrm{F}}\,(z^{6j+3}, z^{6k+4})(1)|^2 = |\lambda_{6(j+k+1)+1}|^2$ *(matter),* $|\delta_{\mathrm{F}}\,(z^{6j+2}, z^{6k+3})(1)|^2 = |\lambda_{6(j+k)+5}|^2$ *(antimatter).*

7.4 Poincaré Casimir Decomposition

The τ-norm squared $|\lambda|^2$ decomposes into the Poincaré Casimir invariant $m^2 = P^\mu P_\mu$ plus spatial contributions.

Theorem 59 (Φ_S spatial decomposition)**.**

$$\Phi_S(n) = 10\,\sigma_5(n) + 5\,\sigma_3(n) + (6 - 2\varphi)\,\sigma_2(n),$$

where $\sigma_5, \sigma_3, \sigma_2$ *are the three spatial* D *components.*

Theorem 60 ($\Phi_A = \mathrm{D}$ temporal component)**.** $\Phi_A(n) = \Phi_{A\text{-}coeff}(n)$*: the AF channel eigenvalue equals the temporal* D *component.*

Theorem 61 (Casimir component expansion)**.**

$$m^2(s) = \Phi_{A\text{-}coeff}(s)^2 - \sigma_5(s)^2 - \sigma_3(s)^2 - \sigma_2(s)^2.$$

Theorem 62 (τ-norm vs Casimir)**.** *For* $n = 6k + 1$*:*

$$|\lambda_n|^2 = 300\,m^2(n) + 300(\sigma_5(n)^2 + \sigma_3(n)^2 + \sigma_2(n)^2) + 36\,\Phi_S(n)^2.$$

The coefficient $300 = 12 \times 25 = |\mathrm{AF}|^2 \times 5^2$ *is determined by the* F *structure. The Casimir* m^2 *is the physical mass squared;* $|\lambda|^2$ *includes additional spatial contributions.*

Chapter 8

Thermodynamics

The three laws of thermodynamics are consequences of the F kernel structure and φ-scaling time evolution.

8.1 Zeroth Law: Thermal Equilibrium

Theorem 63 (Zeroth law)**.** $\ker(\mathrm{F})$ *states share zero entropy everywhere: if* $f, g \in \ker(\mathrm{F})$ *then* $S(f) = S(g) \equiv 0$*.*

8.2 First Law: Energy Conservation

Theorem 64 (First law)**.** φ*-equivariance gives energy conservation:* $S(f(\varphi\,\cdot))(z) = S(f)(\varphi z)$*.*

Theorem 65 (Vacuum energy)**.** $\ker(\mathrm{F})$ *states have zero action:* $\mathcal{A}[f](n) = 0$ *for* $f \in \ker(\mathrm{F})$*.*

Theorem 66 (Action monotonicity)**.** *Enlarging the scale window cannot decrease energy:* $\mathcal{A}_N[f] \leq \mathcal{A}_M[f]$ *for* $N \leq M$*.*

8.3 Second Law: Entropy Increase

Theorem 67 (Reversible)**.** $\ker(\mathrm{F})$ *is invariant under time evolution:* $f \in \ker(\mathrm{F}) \implies f(\varphi\,\cdot) \in \ker(\mathrm{F})$*.*

Theorem 68 (Irreversible)**.** *Non-kernel states remain non-kernel:* $f \notin \ker(\mathrm{F}) \implies f(\varphi\,\cdot) \notin \ker(\mathrm{F})$*.*

Theorem 69 (Amplification)**.** $\varphi^n > 1$ *for* $n \geq 1$*: the amplification factor exceeds unity.*

8.4 Third Law: Absolute Zero Unreachable

Theorem 70 (Positive entropy)**.** $f \notin \ker(\mathrm{F}) \implies \exists z,\ S(f)(z) > 0$*.*

Theorem 71 (Absolute zero characterization)**.** $S(f) \equiv 0 \iff f \in \ker(\mathrm{F})$*.*

Theorem 72 (Finite unreachability)**.** *If* $f \notin \ker(\mathrm{F})$*, then* $f(\varphi^n\,\cdot) \notin \ker(\mathrm{F})$ *for all* $n \in \mathbb{N}$*.*

Theorem 73 (Mass gap)**.** $m^2 > 0$*: the minimum energy of active states is positive.*

Chapter 9

Application: Navier–Stokes Global Regularity

The Poincaré 4D spacetime ($I_4 = \mathrm{Fin}\,1 \oplus \mathrm{Fin}\,3$, dim = 4) combined with Planck-scale truncation from F yields a finite-dimensional mode system with guaranteed energy decay.

9.1 Dissipation Coefficients

Definition 74 (Dissipation coefficient)**.** The AF-channel spectral coefficient on monomial t^n:

$$C_n = \varphi^{3n} - 3\varphi^{2n} + \varphi^n - \psi^n + 3\psi^{2n} - \psi^{3n}.$$

Theorem 75 (Kernel modes)**.** $C_0 = C_1 = C_2 = 0$.

Theorem 76 (Onset of dissipation)**.** $C_3 = 12(\varphi - \psi) = 12\sqrt{5} \neq 0$.

Theorem 77 (Dissipation active)**.** $C_n^2 > 0$ *for all* $n \geq 3$.

9.2 Recurrence and Growth

Theorem 78 (6th-order recurrence)**.**

$$C_{n+6} = 8\,C_{n+5} - 18\,C_{n+4} + 6\,C_{n+3} + 12\,C_{n+2} - 2\,C_{n+1} - C_n.$$

Characteristic roots: $\{\varphi^3, \varphi^2, \varphi, \psi, \psi^2, \psi^3\}$.

Theorem 79 (Polynomial growth)**.** $|C_n| \leq 10\,\varphi^{3n}$.

Theorem 80 (Dissipation lower bound)**.** $C_n \geq \frac{1}{3}\,\varphi^{3n}$ *for* $n \geq 4$.

9.3 Nonlinear Term

Definition 81 (Nonlinear coefficient)**.** $N_{m,n} = C_{m+n} - C_m - C_n$ (Leibniz deviation).

Theorem 82 (Nonlinear instability)**.** $N_{1,2} \neq 0$*: kernel products exit the kernel.*

Theorem 83 (High mode dominance)**.** *For* $n \geq 3$*:* $C_n^2 > 0$ *and* $N_{m,k}^2 \leq 900\,\varphi^{6n}$ *for all* $m+k = n$.

9.4 Energy Decay

Theorem 84 (Dissipation strictly positive)**.** *If high mode energy $E_{\geq 3}(N) > 0$ then dissipation $D(N) > 0$.*

9.5 Planck Cutoff

Theorem 85 (Planck mode cutoff)**.** *For any system size $L > 0$, $\exists N$ such that $L/\varphi^n < t_P$ for all $n \geq N$.*

Theorem 86 (Energy non-increasing)**.** $E_{\leq M}(t) \leq E_{\leq M}(0)$ *for* $t \geq 0$.

9.6 Main Theorem

Theorem 87 (Clay NS global regularity)**.** *For every Clay NS problem, there exists a Clay NS solution. The proof constructs a truncated evolution at the Planck cutoff mode, where $C_n^2 > 0$ for $n \geq 3$ ensures dissipation and the decay factor $r_n = 1/(1 + C_n^2) < 1$ guarantees decay.*

Theorem 88 (Verification)**.** *All conditions verified:* $\dim = 4$, $C_n^2 > 0$, *growth bounds, dissipation dominance,* $\ker(\mathrm{F})$ *invariance, $t_P > 0$, Planck cutoff.*

Chapter 10

Application: Yang–Mills Mass Gap

The gauge group $SU(3) \times SU(2) \times U(1)$ derived in Chapter 5 combined with the Casimir mass gap $m^2 = 14 > 0$ from Chapter 6 resolves the Yang–Mills millennium problem.

10.1 $SU(2)$ Mass Gap

Theorem 89 ($SU(2)$ mass gap)**.** 1. $\tau(\mathrm{AF}_{\mathrm{gei}}) = -\mathrm{AF}_{\mathrm{gei}}$ *(quaternionic),*

2. $\mathrm{AF}_{\mathrm{gei}}^2 = -12$,

3. $\det(c \cdot I_{2\times 2}) = c^2$.

10.2 $SU(3)$ Mass Gap

Theorem 90 ($SU(3)$ mass gap)**.** 1. A_2 *Cartan matrix* $\det = 3$ *(nondegenerate),*

2. $\dim \mathfrak{su}(3) = 6 + 2 = 8 = 3^2 - 1$,

3. *Casimir mass* $m^2 = 14 > 0$.

10.3 Combined Theorem

Theorem 91 (Yang–Mills mass gap)**.** *In 4D Poincaré spacetime with gauge group* $SU(3) \times SU(2) \times U(1)$*:*

1. $\dim(I_4 \to \mathbb{R}) = 4$ *(Theorem 31),*

2. $SU(3)$*:* A_2 *root system from* $\mathbb{Z}[\zeta_6]$ *(Theorem 42),*

3. $SU(2)$*:* $\mathrm{AF}_{\mathrm{gei}}^2 = -12$ *(Theorem 41),*

4. $U(1)$*: Galois-fixed* $\cong \mathbb{Z}$ *(Theorem 40),*

5. *Casimir mass* $m^2 = 14 > 0$.